

Schwarzschild-De Sitter black holes in 4 + 1 dimensional bulk

*Metin ARIK and †Dilek ÇİFTÇİ

Department of Physics, Boğaziçi University, 34342 Bebek, Istanbul TURKEY

February 7, 2008

Abstract

We construct a static solution for 4+1 dimensional bulk such that the 3+1 dimensional world has a linear warp factor and describes the Schwarzschild-dS₄ black hole. For $m = 0$ this four dimensional universe and Friedmann–Robertson–Walker universe are related with an explicit coordinate transformation. We emphasize that for linear warp factors the effect of bulk on the brane world shows up as the dS₄ background which is favored by the big bang cosmology.

Keywords: Schwarzschild-dS₄ black hole; Randall–Sundrum models; Brane world; Hubble length.

In the universe, there exists four fundamental interactions; weak, strong, electromagnetic and gravitational. However, all the interactions in nature are not unified. Gravity shows some different aspects because of its long range and unquantizable behavior and its importance in cosmology. To get a natural explanation for the unification problem, the possibility that gravity might not be fundamentally four dimensional has been considered [1]-[4]. Basically these scenarios say that our universe may be a brane embedded in some higher dimensional space. All matter and gauge interactions live on the brane while gravitational interactions are effective in the whole higher dimensional space. Besides cosmology, black holes explore the geometry of gravity and its quantum effects. Therefore gravity and black holes can be related with the brane-world cosmology. In many of the brane-world scenarios, the matter fields we observe are trapped on the brane and when they undergo gravitational collapse a black hole forms [5]-[16]. This black hole becomes a higher dimensional object. This higher dimensional object can recover the usual physical properties of black holes and stars.

Most of the brane world scenarios originated from Randall–Sundrum (R-S) models in which the four dimensional Minkowski universe is embedded in five dimensional AdS bulk. This bulk is compatible with a Friedmann–Robertson–Walker (FRW) brane. The most general vacuum bulk with a FRW brane is Schwarzschild-anti-de Sitter space-time [17],[19] where a domain wall moving in the 5 dimensional Schwarzschild-AdS space-time with the metric

$$\begin{aligned} ds^2 &= g_{\mu\nu} dx^\mu dx^\nu \\ &= -f(r) dt^2 + \frac{dr^2}{f(r)} + r^2 [d\chi^2 + f_k(\chi) d\Omega_2^2] \end{aligned} \quad (1)$$

is considered. Here the part in the square parenthesis is the metric of a unit three-dimensional sphere, plane or hyperboloid for $k = 1, 0, -1$ and $f_1(\chi) = \sin \chi$, $f_0(\chi) = \chi$, and $f_{-1}(\chi) = \sinh \chi$ respectively. The solution of field equations $R_{\mu\nu}^5 = \Lambda^5 g_{\mu\nu}^5$ with negative cosmological constant $\Lambda^5 = -\frac{4}{l^2}$ is given by

$$f(r) = k + \frac{r^2}{l^2} - \frac{\mu}{r^2},$$

where l and μ are constants. For $\mu = 0$, this reduces to AdS₅ and $\mu \neq 0$ generates the electric part of the Weyl Tensor. By this consideration the location of three branes in the five dimensional bulk can be investigated.

Instead of these theories we have considered a four dimensional curved space-time with positive cosmological constant embedded in five dimensional flat and empty Minkowski Universe as shown in [20]. Starting from

*metin.arik@boun.edu.tr

†dciftci@boun.edu.tr

this point of view, in this paper, we consider a usual four dimensional Schwarzschild metric embedded in five dimensional Ricci flat universe. Our metric describes a black hole on a domain wall at fixed w . As such a space we take the five dimensional analogue of Schwarzschild-de Sitter space-time in the following form.

$$ds^2 = dw^2 + \left(1 - \frac{|w|}{w_0}\right)^2 \left[-U(r) dt^2 + \frac{1}{U(r)} dr^2 + r^2 d\Omega_2^2\right], \quad (2)$$

where we choose

$$U(r) = 1 - \frac{2M}{r} - \frac{r^2}{w_0^2}, \quad (3)$$

such that for $M = 0$ the bulk is flat and for $M \neq 0$ the bulk is Ricci flat. This metric tells us that for constant w the cosmological constant is $\Lambda_{observed} = 3/w_0^2$. It is precisely the condition that the parameter w_0 in $U(r)$ Eq. (3) and in the warp factor in Eq. (2) are the same which causes the 4 + 1 dimensional bulk to be Ricci flat. We set up the 3 + 1 branes at $w = 0$ and $w = w_1 \leq w_0$. Then, on the branes this metric will be in the standard form of the Schwarzschild metric. Here note that in small scales the cosmological curvature is much smaller than the Schwarzschild curvature. If we denote the perturbed horizon as $r = 2M + \varepsilon$ for small ε , the black hole horizon is determined to be at $r = 2M + 8M^3/w_0^2$ where $w_0 \gg M$. On the other hand from the square of the Riemann tensor one finds curvature singularities at $r = 0$ and $w = \pm w_0$. However one expects that masses confined to the brane will generate the localized gravitational fields, so that these singularities are not physically important. Strictly speaking the warp factor is given by $1 - |w|/w_0$, for $-w_1 \leq w \leq w_1$, and imposing periodicity of period $2w_1$ together with Z_2 symmetry $w \rightarrow -w$.

For $M = 0$ the four dimensional part in square parenthesis in Eq. (2) transforms into

$$ds_4^2 = \left[-d\tau^2 + g(\tau)^2 \left(d\chi^2 + f_k(\chi)^2 d\Omega_2^2\right)\right], \quad (4)$$

by the following transformations. Here τ and $g(\tau)$ correspond to cosmic time and scale factors of FRW universe respectively.

- $k = -1$

$$\begin{aligned} r &= w_0 \sinh(\tau) \sinh(\chi) \\ t &= \frac{w_0}{2} \ln \left(\frac{\cosh(\chi) \sinh(\tau) + \cosh(\tau)}{\cosh(\chi) \sinh(\tau) - \cosh(\tau)} \right) \\ g(\tau) &= w_0 \sinh(\tau) \\ f_{-1}(\chi) &= \sinh(\chi). \end{aligned}$$

- $k = 0$

$$\begin{aligned} r &= \chi \exp(\tau/w_0) \\ t &= \tau - \frac{w_0}{2} [\chi^2 \exp(2\tau/w_0) - 1] \\ g(\tau) &= \exp(\tau/w_0) \\ f_0(\chi) &= \chi. \end{aligned}$$

- $k = 1$

$$\begin{aligned} r &= w_0 \cosh(\tau) \sin(\chi) \\ t &= \frac{w_0}{2} \ln \left(\frac{\cosh(\tau) \cos(\chi) + \sinh(\tau)}{\cosh(\tau) \cos(\chi) - \sinh(\tau)} \right) \\ g(\tau) &= w_0 \cosh(\tau) \\ f_1(\chi) &= \sin(\chi). \end{aligned}$$

For the case of $M \neq 0$, we can make the same transformation, however it will not make our work easier. If we put $M = 0$ in this transformation we obtain the same metric as in Eq. (4). This means that, our space-time becomes FRW universe far away from the black hole. Hence we can use the context of FRW cosmological model. In this part we suppose that the visible world is located at $w = w_1$. Therefore the visible metric tensor will be $g_{\mu\nu}^{(vis)} = g_{\mu\nu}(x^\mu, w = w_1)$ and for the hidden brane $g_{\mu\nu}^{(hid)} = g_{\mu\nu}(x^\mu, w = 0)$. The setup is similar to the scenario of [2]. At first, in order to get a realistic theory we need to look for the localization of gravity on the brane. To do this, taking $(1 - |w|/w_0)g_{\mu\nu} + h_{\mu\nu}(x, w)$, we study the fluctuations of the metric. Here $h_{\mu\nu}(x, w)$ is the perturbed metric and its transverse traceless components represents the graviton on the brane i.e. the conditions $\partial_\mu h^{\mu\nu} = 0$ and $h^\mu_\mu = 0$ must be satisfied [1],[2],[21]-[25]. By separation of variables we write $h_{\mu\nu}$ in terms of the four-dimensional mass eigenstates $h_{\mu\nu}(x, w) = \phi_{\mu\nu}(t, x^i)\psi(m, w)$, where m corresponds to four-dimensional mass. Performing a coordinate transformation from w to the variable z defined by $\partial z / \partial w = \pm (1 - |w|/w_0)^{-1}$ and defining $u(z)$ by $\psi = (1 - |w|/w_0)^{-3/2} u(z)$ we get the wave function as the solution of

$$(-\partial_z^2 + V(z))u(z) = m^2 u(z), \quad (5)$$

where the potential $V(z)$ is

$$V(z) = \frac{3}{4} \frac{f'^2}{f^2} + \frac{3}{2} \frac{f''}{f}. \quad (6)$$

Here we denote the derivative with respect to z by prime and f is the z dependent warp factor as addressed below. Since our brane is located at $w = w_1$, we can write the warp factor as $f(w) = 1 - \frac{w_1}{w_0} + \frac{|w_1 - w|}{w_0}$, then z dependence of the warp factor will be

$$f(z) = \left(1 - \frac{w_1}{w_0}\right)^{-1} \exp(-|z|/w_0), \quad (7)$$

the transformed coordinate z is given by

$$z = \text{sign}(w_1 - w) \ln \left(\frac{1 - \frac{w_1}{w_0}}{1 - \frac{w_1}{w_0} + \frac{|w_1 - w|}{w_0}} \right), \quad (8)$$

where our brane is placed at $z = 0$. Then we find the potential as

$$V(z) = \frac{9}{4w_0^2} - \frac{3}{w_0} \delta(z). \quad (9)$$

Here the δ function in the potential allows a normalizable bound state mode and this shows that the 5D graviton is localized on the brane. In general for a potential $V(z)$ which allows bound states, Eq. (5) has $m = 0$ normalizable solutions iff $V(z) > 0$, as $z \rightarrow \infty$. In this case localization of gravity on the brane occurs. Here $m = 0$ mode corresponds to the gravitons on the brane.

The next point is to solve the Schrodinger Equation (5). First of all, for the zero mode wave function, we find that $u_0(z) = (3/2w_0)^{1/2} \exp(-3w_0|z|/2)$. On the other hand, the massive modes are $u_m(z) = A \exp\left(-\sqrt{\frac{9}{4w_0^2} - m^2}|z|\right)$, where A is the normalization factor. As explained in [24], the massive modes for $0 < m^2 < 9/4w_0^2$ do not exist there and for $m^2 \geq 9/4w_0^2$ this wave function becomes plane wave. Then there is a mass gap $9/4w_0^2$ between the zero mode and the continuous modes.

At this stage, we look for the five dimensional Einstein tensor,

$$G_{MN}^5 = -3\delta_M^\mu \delta_N^\nu \eta_{\mu\nu} \frac{1}{(w_0 - |w|)} [\delta(w) - \delta(w - w_1)], \quad (10)$$

where $\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$, Capital Latin letters $M, N = 0, 1, 2, 3, 5$ denote full space-time and lower Greek $\mu, \nu = 0, 1, 2, 3$ run over the brane world. From Eq. (10), it is easily seen that there are no nongravitational fields ($T_{MN} = 0$) and no cosmological constant in the bulk. For the solution on the brane, we introduce the boundary terms as

$$V_{vis} = \frac{-24M_5^3}{w_0 \left(1 - \frac{|w_1|}{w_0}\right)} \quad \text{and} \quad V_{hid} = \frac{24M_5^3}{w_0}. \quad (11)$$

These results are similar but not the same as RS model in which V_{vis} and V_{hid} are same with opposite sign. We can derive the induced brane gravity [8], which consists of four dimensional Einstein gravity on the brane. From the dimensionful constants κ_5 , and κ_4 the Planck masses M_5 , M_4 are defined as

$$\kappa_5^2 = 8\pi G_{(5)} = M_5^{-3}, \quad \kappa_4^2 = 8\pi G_{(4)} = M_4^{-2}. \quad (12)$$

The induced Einstein tensor on the brane is

$$\frac{M_4^2}{2} \int d^4x \sqrt{g^4} G_{\mu\nu}^4 = -\frac{6M_5^3}{2} \int d^4x \int_{-w_1}^{w_1} \frac{dw}{w_0} \sqrt{g^5} g_{\mu\nu} \left(1 - \frac{|w|}{w_0}\right)^{-1} [\delta(w) - \delta(w - w_1)], \quad (13)$$

then for the hidden brane world

$$G_{\mu\nu}^4(hid) = -\frac{3}{w_0^2} \eta_{\mu\nu} \quad \text{and} \quad M_4^2 = \frac{V_{hid} w_0^2}{12}, \quad (14)$$

and for the visible world

$$G_{\mu\nu}^4(vis) = -\frac{3}{w_0^2 \left(1 - \frac{w_1}{w_0}\right)^2} \eta_{\mu\nu} \quad \text{and} \quad M_4^2 = \frac{V_{vis} w_0^2}{12} \left(1 - \frac{w_1}{w_0}\right)^2, \quad (15)$$

where M_4 and M_5 have dimensions (length) $^{-1}$ [8] and w_0 corresponds to the $1/k$ term in [2]. The Einstein tensor on the brane is found as exactly what we want to get. From the four dimensional part in the square parenthesis in Eq. (2) we determine the same result for the Einstein tensor or matter content. This is the usual well known Schwarzschild space-time on the visible and hidden world. Therefore we get that the cosmological constant of the hidden world is $\Lambda_{4(hid)} = 3/w_0^2$. On the other hand for the visible brane we get the positive cosmological constant $\Lambda_{4(vis)} = 3(w_0 - w_1)^{-2}$ and the five dimensional cosmological constant Λ_5 is zero everywhere. This may be interpreted as the four dimensional cosmological constant originating from the localized energy momentum tensor on the hidden brane affecting the visible brane through the bulk. In Eq. (11) the minus sign on the visible brane tension comes from the attractive gravitational force. This minus sign causes us to measure positive cosmological constant on the visible world.

To find the mass parameter on the 3-brane we find the effective action as in [2]. Then we get

$$m = \left(1 - \frac{|w_1|}{w_0}\right) m_0. \quad (16)$$

Here if the part in parenthesis is of order 10^{-15} , near the curvature singularity $w_1 \simeq w_0$ where the visible world is located, we get the TeV mass scale from the Planck mass scale, $10^{16} TeV$.

These results have some importance in some sense:

- $V_{vis} = -10^{15} V_{hid}$.
- The observed cosmological constant ($10^{-52} m^{-2}$) comes from the effect of the hidden brane on our world. Really what we observe as a cosmological constant $10^{-52} m^{-2}$ is the $\Lambda_{4(hid)}$. From this consideration we find w_o from the equation

$$10^{-52} m^{-2} = \frac{3}{w_0^2}, \quad (17)$$

then $w_1 \simeq w_o = 10^{26} m$ is approximately just the Hubble length. Therefore we are located at the distance of nearly Hubble length from the hidden brane. On the other hand, the cosmological constant of the visible world which we cannot observe is $10^{-22} m^{-2}$. The factor 10^{30} in the two cosmological constants comes from the location of our world. We are much closer to singularity w_0 than the hidden brane. Really there is no cosmological constant more than one. To prevent this complication, we do not use the name $\Lambda_{(vis)}$ anymore, we call it $G_{00}(w)$. We can summarize these results, by the relation

$$\Lambda_{(hid)} = \Lambda = G_{00}(w) \left(1 - \frac{|w|}{w_0}\right)^2 = 10^{-52} m^{-2}, \quad (18)$$

which is satisfied for all w . Placing our brane at $w = w_1$ causes $G_{00}(w_1)$ to be $10^{-22}m^{-2}$. If we had placed our brane at any other w , G_{00} on the brane would change as indicated in this equation but Λ would not change. $G_{00}(w_0)$ becomes infinite on the singularity at $w = w_0$. However $0 \leq w \leq w_1 < w_0$ and the point $w = w_0$ is outside the branes and the bulk.

One of the main motivations for this work has been the argument that, since there is nothing except gravity in the fifth dimension, there should also be no cosmological constant in the fifth dimension. Gravity originates from the hidden brane and then it is localized and interacts with the matter on the brane we live. We cannot see the hidden brane. However we can measure its effect as a cosmological constant Λ . Additionally the distance from the hidden brane determines the localized mass density of the universe. Namely $G_{00}(w_1)$, the mass density in the visible brane depends on Λ and varies with the location of the universe in the bulk. This causes the energy density to be obtained as $\rho \simeq 10^3 kg/m^3$. This value is too large to be interpreted as an average energy density for our universe. However, interestingly enough this value is comparable to stellar energy densities. This indicates that a model where the cosmological constant is localized and creates localized masses may be viable.

At the end, the correction to Newton's Law is obtained as

$$\begin{aligned} U(r) &= -G_N \frac{m_1 m_2}{r} - M^{-3} \int_{m_0}^{\infty} dm \frac{m_1 m_2 e^{-mr}}{r} (u_m(0))^2 \\ &= -G_N \frac{m_1 m_2}{r} (1 + C \frac{e^{-m_0 r}}{r}), \end{aligned} \quad (19)$$

where as we have explained above $m_0 = 3/2w_0$ and $A^2 = \left[\ln \left(1 - \frac{w_1}{w_0} \right)^{-1} \right]^{-1}$, and then the constant $C = 16\pi w_0 \left(1 - \frac{w_1}{w_0} \right) \left(-\ln \left(1 - \frac{w_1}{w_0} \right) \right)^{-1} \simeq 10^{11}m$. This value of C is severely in violation of experimental measurements.

In conclusion, in this paper, we have presented a new brane world black hole solution such that the Schwarzschild-dS₄ space-time is embedded into 4 + 1 dimensional bulk with the linear warp factor $(1 - |w|/w_0)$. Here the visible brane is located at $w = w_1 < w_0$. We have presented the coordinate transformation between the Schwarzschild-dS₄ and FRW space-time for $M = 0$. The same transformation can also be used for $M \neq 0$. Although such a transformation will result in a complicated metric, for $M = 0$ this metric will reduce to FRW. For $M \neq 0$, we found that in the bulk not only the energy momentum tensor but also the cosmological constant is zero. However the effect of the hidden brane to our world is just the cosmological constant which we now measure. The localized energy momentum tensor in the bulk also serves to cause a four dimensional cosmological constant on the brane. It was found that this cosmological constant depends on the distance scale w_0 . Finally we have derived the range of fifth dimension from the hierarchy between the weak and Planck scales. This model is in violation of observed values concerning corrections to the Newtonian corrections and the matter-average energy density of the universe.

References

- [1] L. Randall, R. Sundrum, *Phys. Rev. Lett.* **83**, 4690-4693 (1999).
- [2] L. Randall, R. Sundrum, *Phys. Rev. Lett.* **83**, 3370-3373 (1999).
- [3] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, *Phys. Lett.* **B436**, 257 (1998).
- [4] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, *Phys. Rev.* **D59**, 086004 (1999).
- [5] A. Chamblin, H.S. Reall, H.A. Shinkai, and T. Shiromizu, *Phys. Rev.* **D63**, 064015 (2001).
- [6] A. Chamblin, S.W. Hawking, and H.S. Reall, *Phys. Rev.* **D61**, 065007 (2000).
- [7] R. Gregory, R. Whisker, K. Beckwith, and C. Done, *JCAP* **10**, 013 (2004).
- [8] G. Kofinas, *Phys. Rev.* **D66**, 104028 (2002).

- [9] Y. S. Myung, *Class. Quantum Grav.* **21**, 1279 (2004).
- [10] S. Nojiri, S. D. Odinstov, S. Ogushi, *Phys. Rev.* **D65**, 023521 (2002).
- [11] N.K. Dadhich, R. Maartens, P. Papadopoulos, and V. Rezanian, *Phys. Lett.* **B487**, 1 (2000).
- [12] R. Maartens, *Living Rev. Relativity* **7**, (2004).
- [13] H. Takayuki, and K. Gungwon, *Phys. Rev.* **D64**, 064010 (2001).
- [14] R. Casadio, A. Fabbri, and L. Mazzacurati, *Phys. Rev.* **D65**, 084040 (2002).
- [15] R. Casadio, L. Mazzacurati, *Mod. Phys. Lett.* **A18**, 651 (2003).
- [16] R. Emparan, G.T. Horowitz, and R.C. Myers, *JHEP* **01**, 021 (2000.)
- [17] D. Ida, *JHEP* **09**, 014 (2000).
- [18] S. Mukohyama, T. Shiromizu, and K. Maeda, *Phys. Rev.* **D62**, 024028 (2000).
- [19] P. Bewcock, C. Charmousis, and R. Gregory, *Class. Quantum Grav.* **17**, 4745 (2000).
- [20] M. Arık, et al., *Phys. Rev.* **D68**, 123503 (2003).
- [21] I. Brevik, K. Ghoroku, S. D. Odintsov, M. Yahiro, *Phys. Rev.* **D66**, 064016 (2002).
- [22] C. Csaki, J. Erlich, T. J. Hollowood, Y. Shirman, *Nucl. Phys.* **B581**, 309 (2000).
- [23] J. F. Vazquez-Poritz, *JHEP* **0209**, 001 (2002.)
- [24] M. Ito, *Europhys. Lett.* **64**, 295 (2003).
- [25] T. Padmanabhan, S. Shankaranarayanan, *Phys. Rev.* **D63**, 105021 (2001).
- [26] J. Garriga, T. Tanaka, *Phys. Rev. Lett.* **84**, 2778 (2000).